

Insulin adsorption on different Transfersomes




Fig. 1

e.g. examples 1-27, A

Insulin adsorption on Transfersomes C




Fig. 2

e.g. examples 1-27, B

Insulin adsorption on different Transfersomes




Fig. 3

e.g example 1-27, C

Insulin adsorption on different Transfersomes




Fig. 4

e.g example 46-59

Insulin adsorption to different Transfersomes




Fig. 5

examples 72-76

Insulin adsorption on different Transf rsomes




Fig. 6

example 77-92

Transfersomes comprising
SPC+SPG/Tween = L/D = 2/1




Fig. 7

examples 96-98



Insulin association with
Transfersomes C

Fig. 8

examples 99-100

insulin association with TransfersomesTM

- A: SPC/NaCholate in cholate buffer + ActrapidTM
 B: SPC/NaCholate, 5 % + Actrapid
 C: SPC/NaCholate, 10 % + Actrapid
 D: SPC/SPG/Tween80 (3/1/1) + Actrapid
 E: SPC/NaCholate + lyophilized human insulin in buffer
 F: SPC/NaCholate + Velasulin (porcine insulin)
 G: SPC/Tween 80 (2/1) + Actrapid, incubated for 5 weeks
 H: SPC/Tween 80 (2/1) + Actrapid, incubated for 4 days
 I: SPC/Tween 80 (2/1) + Actrapid, incubated for 3 hours
 K: SPC/Tween 80 (2/1) + Actrapid, incubated for 2 hours
 L: SPC (liposomes), 10 % stock susp.

Fig. 9

selected, representative, results




Figure 10

004T30" 33655510




Figure 11

epicutaneous
Transfersulin C, 40 IU

Insulin batch effect




Figure 12